شركة مام ايجيبت للأدوية

als

MAMIZOLE 12% Injectable solution السيانات العلمية مطابق كما تم الموا فق ملي من قبل اللجن العلمية المدوية السطر و المانات (For vet use only) الاعلاف دا مسرياع ما هر

Composition:

Each100 ml contains:

17.1 gm of imidocarp dipropionate (eq. to 12 gm imidocarb base)

Properities:

Imidocarb dipropionate is a substituted carbinilide, used as an antiprotozoan treatment for the control of babesia spp. Little is known about the mode –of- action of imidocarb dipropionate . it appears that imidocarb acts directly on the parasite, causing alteration in number and size of nuclei and in morphology(vacuolation) of the cytoplasm . the antiprotozoan activity is derived from the carbanilide acting on glycolyis of the parasite . this is the result of this class of drugs giving rise to hypoglycaemia in the host. Babesia as well as many other parasites like trypanosomes depends upon host glucose for aerobic glycolysis. There is also a selective blocking effect on the replication of the quinetoplastic DNA of the parasite

Indication

For treatment and prevention of bovine (red water fever - babesia divergens infection) only.

Target spp:.

Cattle.

Dosage & administration

- -Route of administration subcutaneous injection only
- -Dose of whole product

For treatment: 0.70 ml / 100 kg body weight once.

For prevention: 1.80 ml/ 100 kg body weight once.

Warnings and precautions:

Not for use in any other species.

- -Special precautions for use in animals
- -Estimate bodyweight carefully and do not exceed the recommended dosage
- -Special precautions for the person administering the veterinary medicinal product to animals
- -Do not use if under medical advice not to work with compounds which may exhibit anti-
- -Wash splashes of the product off the skin and eyes immediately.

Wear suitable protective clothing(ie impermeable gloves) when using the product .

may no

المبيعات والمخازن : شبين الكوم :

۸۷ ش مصطفی کامل البحری البر الشرقی

موبایل : ۱۰۸۸۳۲۵۹۰ فاکس : ۲۷۱۵۹۲۸۸۰۰ الادارة : مدينة العبور - سنتر لؤلؤة العبور -مكتب رقم ٤

موبایل: ۲۰۰۲۰۲۸۳٤٤

E-mail: mam_eg_ph@yahoo.com

شركة مام ايجيبت للأدوية

clay gale

rtra-indications:

Mamizole must not be administerer intravenously or intramuscularly.

-Repeat doses of mamizole must not be given

Adverse effect:

Animals may show cholinergic signs after dosing. It may be possible to alleviate these side effects by treatment with atropine sulphate.

While side-effects (salivation, discomfort, muscle tremors, tachycardia, cough, colics) are rare, they do occur and deaths from anaphylactoid reactions have been recorded following product use.

Withdrawal time:

Cattle: slaughter 213 days, milk 21 days

Storage condition:

Store at a temperature not exceeding 30° C, used 40 days after opening at a temperature not exceeding 30° C.

Pack:

Carton box containing amber glass vial type (II) filled with 100 ml injectable solution with red chlorobutyl rubber stoppers capped with AL/HDPE plastic flip off cap with outer label +insert leaflet.

Manufactured by Arab company for medical products (arabcomed) for Mam Egypt pharmaceuticals



المبيعات والمخازن : شبين الكوم : ۸۷ ش مصطفی کامل البحری البر الشرقی

موبایل : ۱۰۲۸۸۳٤٥٩٠ فاکس: ۲۷۲۰۹۱۸۸۰

الادارة : مدينة العبور - سنتر لؤلؤة العبور -مكتب رقم ٤

موبایل : ۱۰۰۲۰۲۸۳٤٤