

البيانات العلمية مطابقة
للبيانات العلمية المرفقة
بموافقة اللجنة العلمية
د. ياسين عبد السميع

Melomat

Injectable solution

For vet. use only

Composition:

Each ml of Melomat contains 20 mg of meloxicam

Properties:

-Pharmacotherapeutic group:

Anti-inflammatory

-Mode of action:

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, anti-exudative, analgesic and antipyretic properties. Meloxicam also has anti-endotoxic properties because it has been shown to inhibit production of thromboxane B2 induced by E. coli endotoxin administration in calves, lactating cows and pigs

Indications:

in cattle. For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle. For in diarrhea, in combination with oral rehydration therapy, to reduce clinical signs in calves of over one week of age and young non-lactating cattle. For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy. For the relief of post-operative pain following dehorning in calves.

in pigs. For use in noninfectious locomotor disorders to reduce the symptoms of lameness and inflammation. for adjunctive therapy in the treatment of puerperal septicaemia and toxemia (mastitis-metritis-agalactia syndrome) with appropriate antibiotic therapy.

in dogs. For reducing postoperative pain and inflammation following orthopaedic (e.g. fracture operation) and soft-tissue surgery. Moreover, it is used for the alleviation of inflammation and pain in both acute and chronic musculoskeletal disorders in dogs

in cats, for reducing postoperative pain and inflammation after ovariohysterectomy (spay operation), orthopaedic and minor soft-tissue surgery. Moreover, it is used for the alleviation of pain and inflammation in acute and chronic musculoskeletal disorders.

- Scientific Office: 1th floor, 85 Mostafa Kamel El Bahary st.,
Shebin El-Koom, Monofia. Telefax: 048/2194886

- Administration & sales: 7th floor, 85 Mostafa Kamel El Bahary st.,
Shebin El-Koom, Monofia. Telefax: 048/2072153

Website: atm-pharma.com

E-mail: info@atm-pharma.com

د. ياسين عبد السميع

المكتب العلمي ٨٥ مصطفى كامل البحري - الدور الأول علوي - شبن الكوم - المنوفية

تليفون: ٠٤٨٢٠١٩٤٨٨٦

الإدارة والمبيعات ٨٥ مصطفى كامل البحري - الدور السابع - شبن الكوم - المنوفية

تليفون: ٠٤٨/٢٠٧٢١٥٣

Horses:

- For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders.
- For the relief of pain associated with equine colic.

Target species:

Cattle, horses, Pigs, Dogs, Cats

Dosage and administration:

-Route of administration:

Cattle: (SC or IV) , Pigs: (IM) , Horses: (IV)
Dogs: (SC) , Cats: (SC)

-Dosage:

● **Cattle:**

For whole product:

2.5 ml of product / 100 kg b.wt. by single S/C injection or intravenous injection in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

● **Pigs:**

For whole product: 2.0 ml/100 kg b.wt. By single I/M injection in combination with antibiotic therapy, as appropriate. If required, a second administration of meloxicam can be given after 24 hours.

● **Horses:**

For whole product

3.0 ml /100 kg b.wt

Avoid introduction of contamination during use

● **Dogs :**

For whole product: 0.1 ml/10 kg b.wt before surgery by Single subcutaneous injection

● **Cats:**

For whole product: 0.15 ml/10 kg b.wt before surgery by Single subcutaneous injection

Warnings and precautions:

Caution with use in animals less than 6 weeks of age, or aged animals, avoid use in dehydrated, hypovolaemic, hypotensive patients, safety in pregnancy has not been established.

Not for human use

- Scientific Office: 1th floor, 85 Mostafa Kamel El Bahary st.,
Shebin El Koom, Monofia. Telefax: 048/2194886

- Administration & sales: 7th floor, 85 Mostafa Kamel El Bahary st.,
Shebin El Koom, Monofia. Telefax: 048/2072153

Website: atm-pharma.com

E-mail: info@atm-pharma.com

- المكتب العلمي: ٨٥ مصطفى كامل البحري - الدور الأول علوى - شين الكوم - المنوفية

تليفاكس: ٠٤٨/٢١٩٤٨٨٦

- الإدارة والمبيعات: ٨٥ مصطفى كامل البحري - الدور السابع - شين الكوم - المنوفية

تليفاكس: ٠٤٨/٢٠٧٢١٥٣

out of reach of children
not use with newly born pet animals

Contra-indications

Do not use in animal suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastro-intestinal lesions. Do not use in case of hypersensitivity to the active substance or to any of the excipients. For the treatment of diarrhea, do not use in animals less than one week of age.

Side effects

Prolonged use may cause gastro-intestinal lesions, inappetence, vomiting, and diarrhea, rarely transient swelling at injection site.

Withdrawal time:

د. ياسين عبد السميع

Cattle:

Slaughter 15 day, milk: 5 days

Horses: meat and offal: 5 days

Pigs:

Slaughter 5 days.


Storage conditions:

Store at temperature not exceeding 30°C & used after opening within 40 days when stored at temperature not exceeding 30° C

Pack

Carton box containing amber (type II) glass vials filled with 10, 30, 50, 100, 250ml with red chlorobutyle rubber stoppers capped with a flip of aluminum cap with an outer label and insert .

Manufacturer:



Scientific Office: 1th floor, 85 Mostafa Kamel El Bahary st.,
Shebin El-Koom, Monofia. Telefax: 048/2194886

Administration & sales: 3th floor, 85 Mostafa Kamel El Bahary st.,
Shebin El-Koom, Monofia. Telefax: 048/2072153

العنكب القلمى ٨٥ - مصطفي كامل البحرى - الدور الأول عنوى - شبن الكوم - المنوفية
تلفاكس: ٠٤٨/٢١٩٤٨٨٦
الإدارة والمبيعات ٨٥ - مصطفي كامل البحرى - الدور السابع - شبن الكوم - المنوفية
تلفاكس: ٠٤٨/٢٠٧٢١٥٣

Website: atm-pharma.com
E-mail: info@atm-pharma.com